

NATIONAL EXAMINATIONS MAY 2018

04-BS-5 ADVANCED MATHEMATICS

3 Hours duration

NOTES:

1. If doubt exists as to the interpretation of any question, the candidate is urged to submit with the answer paper a clear statement of any assumption made.
2. Candidates may use one of the approved Casio or Sharp calculators. This is a Closed Book Exam. However, candidates are permitted to bring **ONE** aid sheet (8.5"x11") written on both sides.
3. Any five (5) questions constitute a complete paper. Only the first five answers as they appear in your answer book will be marked.
4. All questions are of equal value.

Marking Scheme

1. 20 marks
2. 20 marks
3. (a) 5 marks ; (b) 9 marks ; (c) 6 marks
4. (A) 10 marks ; (B) 10 marks
5. 20 marks
6. (A) (a) 6 marks ; (b) 6 marks; (B) 8 marks
7. (a) 10 marks ; (b) 10 marks

1. Consider the following differential equation

$$(x^2 + 5) \frac{d^2 y}{dx^2} + 3x \frac{dy}{dx} + 3y = 0$$

Find two linearly independent solutions about the ordinary point  $x=0$ .

2. Find the Fourier series expansion of the periodic function  $f(x)$  of period  $p = 2\pi$ .

$$f(x) = \begin{cases} \frac{\pi}{2} & -\pi < x \leq -\frac{\pi}{2} \\ x + \pi & -\frac{\pi}{2} < x < 0 \\ \pi - x & 0 \leq x < \frac{\pi}{2} \\ \frac{\pi}{2} & \frac{\pi}{2} \leq x < \pi \end{cases}$$

3. Consider the following function where  $a$  is a positive constant

$$f(x) = \begin{cases} \frac{a}{4} \cos(ax) & -\frac{\pi}{2a} < x < \frac{\pi}{2a} \\ 0 & \text{otherwise} \end{cases}$$

(a) Compute the area bounded by  $f(x)$  and the  $x$ -axis. Graph  $f(x)$  against  $x$  for  $a = 4$  and  $a = 8$ .

(b) Find the Fourier transform  $F(\omega)$  of  $f(x)$ .

(c) Graph  $F(\omega)$  against  $\omega$  for the same two values of  $a$  mentioned in (a).

Explain what happens to  $f(x)$  and  $F(\omega)$  when  $a$  tends to infinity.

Note: 
$$F(\omega) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) \exp(-i\omega x) dx$$

4.(A) Prove that the coefficients  $\alpha$  and  $\beta$  of the least-squares parabola  $y = \alpha + \beta x^2$  that fits the set of  $n$  points  $(x_i, y_i)$  can be obtained as follows:

$$\alpha = \frac{(\sum_{i=1}^n x_i^4)(\sum_{i=1}^n y_i) - (\sum_{i=1}^n x_i^2)(\sum_{i=1}^n x_i^2 y_i)}{n(\sum_{i=1}^n x_i^4) - (\sum_{i=1}^n x_i^2)^2}; \quad \beta = \frac{n(\sum_{i=1}^n x_i^2 y_i) - (\sum_{i=1}^n x_i^2)(\sum_{i=1}^n y_i)}{n(\sum_{i=1}^n x_i^4) - (\sum_{i=1}^n x_i^2)^2}$$

4.(B) Set up Newton's divided difference formula for the data tabulated below and derive the polynomial of highest possible degree:

x	-4	-3	-2	-1	1	4
F(x)	216	0	-56	-36	16	-56

5. The following results were obtained in a certain experiment:

x	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0	1.5	2.0
y	10.0	63.75	70.0	86.25	80.0	68.75	60.0	61.25	90.0

Use Romberg's algorithm to obtain an approximation of the area bounded by the unknown curve represented by the table and the lines  $x = -2$ ,  $x = 2$  and the  $x$ -axis.

Note: The Romberg algorithm produces a triangular array of numbers, all of which are numerical estimates of the definite integral  $\int_a^b f(x)dx$ . The array is

denoted by the following notation:

$$\begin{matrix} R(1,1) \\ R(2,1) & R(2,2) \\ R(3,1) & R(3,2) & R(3,3) \\ R(4,1) & R(4,2) & R(4,3) & R(4,4) \end{matrix}$$

where

$$R(1,1) = \frac{H_1}{2} [f(a) + f(b)]$$

$$R(k,1) = \frac{1}{2} \left[ R(k-1,1) + H_{k-1} \sum_{n=1}^{n=2^{k-2}} f(a + (2n-1)H_k) \right]; \quad H_k = \frac{b-a}{2^{k-1}}$$

$$R(k, j) = R(k, j-1) + \frac{R(k, j-1) - R(k-1, j-1)}{4^{j-1} - 1}$$

6.(A) (a). The equation  $2e^{-x} - 3\cos x = 0$  has a root between  $a = -1.0$  and  $b = 0$ . Use the method of bisection three times to find a better approximation of this root. (Note: Carry six digits in your calculations).

(b) Use Newton's method twice to find a better approximation of this root. (Note: Carry seven digits in your computations)

6.(B) The equation  $x^4 - 3x^3 + 5 = 0$  has a root in the neighbourhood of  $x_0 = 2$ . Write this equation in the form  $x = g(x)$  and then use fixed-point iteration five times to find a better approximation of this root. (Note: Carry seven digits in your computations)

7. The matrix  $A = \begin{bmatrix} 25 & 5 & 15 \\ 5 & 5 & 9 \\ 15 & 9 & 43 \end{bmatrix}$  can be written as the product  $LL^T$  where

$L = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix}$  and  $L^T$  is the transpose of  $L$ .

(a) Find  $L$ .

(b) Use the results obtained in (a) to solve the following system of three linear equations:

$$25x_1 + 5x_2 + 15x_3 = -10$$

$$5x_1 + 5x_2 + 9x_3 = -1$$

$$15x_1 + 9x_2 + 43x_3 = 8$$

